


Principles for Vaccine Administration at 12 Month Schedule Point



- Administer **Bexsero**[®]
 - ▲ Administer **Prevenar 13**[®]
 - **MMR II**[®] or **Priorix**[®] and **Nimenrix**[®] in either leg
- If administering 2 vaccines at the same site for additional vaccines, separate by 2.5cm

If other vaccines are required at the 12 month schedule point:

1. All vaccines recommended should be given in one visit, if possible
2. If all vaccines recommended cannot be given in one visit, prioritise administration of NIP* recommended vaccines
3. If all NIP vaccines recommended cannot be given, prioritise administration of antigens not previously given
4. If vaccine doses need to be deferred, all outstanding vaccines can be given a minimum of 3 days later
5. Contact the Immunisation Section 1300 232 272 for further advice where the above principles do not apply or do not meet the child's particular situation

Administration of paracetamol with every dose of Bexsero is recommended for children aged less than 2 years.

Never mix vaccines together in the same syringe.

*National Immunisation Program (NIP) Vaccines	Non NIP (State) Vaccines
Nimenrix [®] (IM)	Bexsero [®] (IM) (non-Aboriginal children)
Prevenar 13 [®] (IM)	
MMR II [®] OR Priorix [®] (Subcut/IM)	
Influenza (IM)	
Engerix B [®] paediatric (IM) OR HB Vax II [®] paediatric (IM) (MAR)	
Bexsero [®] (Aboriginal children) (MAR)	

MAR = Medically At Risk
The term Aboriginal is inclusive of Aboriginal and Torres Strait Islander people

For more information:

Immunisation Section
Communicable Disease Control Branch
SA Health
sahealth.sa.gov.au/immunisationprovider



Government of South Australia
SA Health