

After Vaccination Information for Pregnant Women

Vaccines, like any other medication or natural therapy can have side effects. It is important to be aware of any reactions that may occur after receiving the whooping cough (pertussis), flu (influenza) and COVID-19 vaccines.

Are vaccines safe during pregnancy?

Yes. Before a vaccine can be used in Australia it must be licensed by the Therapeutic Goods Administration (TGA). The TGA uses scientific evidence on each vaccine to assess safety and effectiveness. The whooping cough, flu and COVID-19 vaccines have been shown to be safe to have during pregnancy.

Common reactions

Most vaccines can cause mild reactions; these are usually short lasting and do not require special treatment. If the reaction seems severe or persists and/or you are concerned, seek further advice from your GP or immunisation provider as soon as possible or go directly to a hospital. You can also receive general advice by calling:

Immunisation Section at SA Health



1300 232 272



9am to 5pm, Monday to Friday

Healthdirect Australia



1800 022 222



24 hours, 7 days

Rare reactions

On very rare occasions a severe allergic reaction such as anaphylaxis may occur, requiring immediate medical treatment. Your immunisation provider is trained to recognise and manage any immediate severe reactions. A severe anaphylactic reaction will generally occur within the first 15 minutes after receiving a vaccine. Fainting may occur after receiving a vaccine and people should lie down if experiencing lightheadedness.

It is important for you to wait for 15 minutes after receiving a vaccine so you may be observed for any reactions, and for treatment to be provided if needed.

Common reactions	Management of common reactions
All vaccines	
Soreness, redness, itching, swelling or	Paracetamol
burning at the injection site for 1-2 days	Place a cool cloth onto the injection site
Fever (temperature)	Paracetamol
	Have extra fluids
A small hard lump may appear where the	
injection was given and persist for some	No treatment needed
weeks or months	
COVID-19 vaccine	
Fatigue/feeling tired	
Headache	
Fever	Rest
Muscle or joint pain	Paracetamol
Chills/nausea	
Swelling of the lymph nodes	
Diptheria/Tetanus/Pertussis (Whooping Cough) vaccine	
Drowsiness or tiredness	
Feeling unwell, muscle aches and/or	• Rest
joint pain	Paracetamol
Decreased appetite and/or nausea	- Turdectumor
Headache	
Influenza vaccine	
Drowsiness, tiredness and/or feeling unwell	Rest

How to report a reaction

Muscle aches

Reporting a reaction following immunisation is an essential part of ensuring ongoing vaccine safety monitoring. All serious or unexpected reactions should be reported either by completing an online Vaccine Reaction Report Form at www.sahealth.sa.gov.au/immunisation or by contacting the Immunisation Section on 1300 232 272 during business hours. Alternatively you can report the reaction to your immunisation provider.

Paracetamol

Further information

For further information and advice about immunisation and pregnancy, speak to your midwife, doctor or immunisation provider.

Visit the Immunisation and Pregnancy page on the SA Health website at www.sahealth.sa.gov.au/immunisation or National Centre of Immunisation Research and Surveillance at www.ncirs.org.au.

Reference: the Australian Immunisation Handbook at immunisationhandbook.health.gov.au





